



The Impact of Postmortem Interval on the Reliability of Forensic Autopsy Evidence in Murder Cases: A Comprehensive Analysis in Nairobi, Kenya

Stephen Kase^{1*}, Jenne Batdorff¹

¹Department of Forensic Medicine, Agacia Clinic, Nairobi, Kenya

ARTICLE INFO

Keywords:

Evidence reliability
Forensic autopsy
Murder cases
Nairobi
Postmortem interval

***Corresponding author:**

Stephen Kase

E-mail address:

steph.kase@gmail.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.59345/sjfm.v1i1.37>

A B S T R A C T

Introduction: Postmortem interval (PMI) significantly impacts the accuracy and interpretation of forensic autopsy evidence, especially in murder cases. In Nairobi, Kenya, where resources and infrastructure may be limited, understanding the relationship between PMI and evidence reliability is crucial for effective investigations and prosecutions. **Methods:** A retrospective analysis was conducted on 150 murder cases in Nairobi between 2018 and 2023. Cases were stratified based on PMI (early PMI <24 hours, intermediate PMI 24-72 hours, late PMI >72 hours). Autopsy reports were reviewed to assess the reliability of evidence related to the cause of death, manner of death, and identification. Statistical analyses were performed to examine the correlation between PMI and evidence reliability. **Results:** As PMI increased, the reliability of certain forensic evidence decreased significantly. Early PMI cases showed the highest reliability for determining the cause of death, manner of death, and identification. In contrast, late PMI cases often presented challenges in interpreting evidence due to decomposition, scavenging, and environmental factors. This led to uncertainties in determining the cause and manner of death, particularly in cases involving blunt force trauma or asphyxia. **Conclusion:** This study highlights the critical impact of PMI on the reliability of forensic autopsy evidence in murder cases in Nairobi. Early PMI examinations are essential for obtaining accurate and reliable evidence, especially in resource-limited settings. Standardized protocols, improved infrastructure, and ongoing training for forensic pathologists are crucial to mitigate the challenges associated with PMI and ensure justice in murder investigations.

1. Introduction

In the realm of forensic investigations, the pursuit of justice hinges upon the meticulous collection, analysis, and interpretation of evidence. The postmortem examination, or autopsy, stands as a cornerstone of this process, offering critical insights into the cause and manner of death, potential perpetrators, and the circumstances surrounding a crime. The reliability and accuracy of autopsy findings are paramount, not only for successful prosecutions but also for ensuring that the innocent are exonerated and the guilty held accountable. However, the passage

of time between death and autopsy, known as the postmortem interval (PMI), can significantly impact the quality and interpretability of forensic evidence. The PMI acts as an inexorable clock, initiating a cascade of biological, chemical, and environmental processes that progressively alter the deceased's remains. Decomposition, the natural breakdown of tissues and organs, can obscure or even erase evidence of injuries, diseases, or toxins. Scavenging by animals and insects can further complicate the forensic landscape, displacing or destroying vital clues. Environmental factors, such as temperature, humidity, and exposure

to sunlight, can accelerate decomposition or introduce confounding variables. As the PMI lengthens, the challenges facing forensic pathologists escalate, potentially compromising the accuracy and reliability of their findings. In resource-limited settings, such as Nairobi, Kenya, the impact of PMI on forensic investigations is particularly pronounced. Overburdened mortuaries, shortages of trained personnel, and inadequate infrastructure can compound the difficulties associated with PMI. Delayed autopsies, improper storage of remains, and limited access to advanced forensic technologies can further erode the quality of evidence. In such environments, understanding the relationship between PMI and evidence reliability becomes crucial for optimizing investigative strategies and ensuring that justice is served, even in the face of adversity.¹⁻³

The present study seeks to shed light on the complex interplay between PMI and forensic evidence in the context of murder cases in Nairobi. By retrospectively analyzing a cohort of homicide victims, we aim to identify specific challenges associated with PMI and assess its impact on the reliability of autopsy findings. We will focus on key aspects of the investigation, including the determination of cause and manner of death, identification of the deceased, and the detection of potential perpetrators. Accurately determining the cause of death is a fundamental objective of any forensic investigation. However, as PMI increases, the ability to pinpoint the precise cause of death may be compromised. Decomposition can mask or distort injuries, making it difficult to differentiate between antemortem and postmortem trauma. Certain causes of death, such as asphyxia or poisoning, may leave subtle or transient signs that are easily obscured by the ravages of time. In Nairobi, where violent crime is prevalent, understanding the impact of PMI on cause-of-death determination is essential for effective homicide investigations. Distinguishing between natural, accidental, suicidal, and homicidal deaths is a critical task for forensic pathologists. However, as PMI lengthens, the evidence supporting the manner of death may become less clear. Decomposition can mimic or obscure signs of violence, potentially leading to misinterpretations. In

cases where the circumstances surrounding the death are unclear, the impact of PMI on manner-of-death determination can be particularly significant, potentially hindering the pursuit of justice. Identifying the deceased is a crucial step in any forensic investigation, enabling the notification of next of kin and facilitating the grieving process. However, as PMI increases, identification becomes increasingly challenging. Decomposition can distort facial features, fingerprints, and other identifying markers. In cases where the deceased is unknown or lacks readily available identification documents, the impact of PMI on identification can be profound, delaying closure for families and impeding the investigative process.⁴⁻⁷

Forensic evidence plays a pivotal role in identifying and apprehending perpetrators. Trace evidence, such as DNA, fingerprints, or fibers, can link suspects to the crime scene or the victim. However, as PMI increases, the recovery and analysis of trace evidence become more difficult. Decomposition can degrade or destroy biological material, while environmental factors can introduce contamination. In Nairobi, where forensic resources may be limited, understanding the impact of PMI on trace evidence recovery is essential for maximizing the chances of identifying and prosecuting offenders.⁸⁻¹⁰ The present study aims to address these critical issues by examining the impact of PMI on the reliability of forensic autopsy evidence in murder cases in Nairobi, Kenya.

2. Methods

This research employed a retrospective cohort study design to investigate the impact of postmortem interval (PMI) on the reliability of forensic autopsy evidence in murder cases. The study was conducted at the Nairobi City Mortuary, the primary medico-legal facility serving Nairobi, Kenya. This setting was chosen due to its high caseload of homicides, representing a diverse range of causes and manners of death, and its resource-limited environment, reflecting the challenges faced by many forensic institutions in developing countries.

The study population comprised 150 homicide cases autopsied at the Nairobi City Mortuary between January 1st, 2018, and December 31st, 2023. Cases

were selected based on the following inclusion criteria; The victim's death was classified as a homicide; A complete autopsy report was available, including detailed descriptions of external and internal examinations, toxicological analyses, and histological findings; The PMI was recorded in the autopsy report; Cases with extensive decomposition or skeletalization were excluded to focus on the impact of PMI within the early postmortem period, where changes are more dynamic and impactful on evidence interpretation. Data were extracted from autopsy reports using a standardized data collection form. The following variables were recorded; Victim demographics: Age, sex, ethnicity; Case characteristics: Date and time of death, date and time of autopsy, location of death, circumstances surrounding the death (as documented in the police report or witness statements); PMI: Recorded in hours based on the pathologist's estimation, considering factors such as body temperature, rigor mortis, livor mortis, and scene investigation findings; Cause of death: As determined by the forensic pathologist, categorized according to the International Classification of Diseases (ICD-10); Manner of death: Classified as natural, accidental, suicidal, or homicidal; Identification method: Documented method used to identify the deceased, such as visual recognition, fingerprints, dental records, or DNA analysis; Evidence reliability: Assessed for each case based on the clarity and interpretability of autopsy findings related to cause of death, manner of death, and identification. Findings were clear, unambiguous, and supported by multiple lines of evidence. Findings were suggestive but not conclusive, or some ambiguity existed due to decomposition or other factors. Findings were inconclusive or significantly obscured by postmortem changes, making definitive interpretations challenging. Data collection was performed by two trained research assistants who underwent a rigorous training program to ensure consistency and accuracy in data extraction. A third researcher independently reviewed a random sample of 20% of the cases to assess inter-rater reliability.

Data were entered into a secure database and analyzed using SPSS version 27.0. Descriptive

statistics were used to summarize the characteristics of the study population and the distribution of PMI, cause of death, manner of death, and identification methods. The impact of PMI on evidence reliability was assessed using the following approaches; PMI categorization: PMI was categorized into three groups; Early PMI: <24 hours; Intermediate PMI: 24-72 hours; Late PMI: >72 hours. Chi-square tests of independence were performed to examine the association between PMI categories and the reliability of evidence related to cause of death, manner of death, and identification. Logistic regression models were developed to assess the independent effect of PMI on evidence reliability, adjusting for potential confounders such as victim demographics, cause of death, and manner of death. Sensitivity analyses were conducted to assess the robustness of the findings to variations in PMI categorization and evidence reliability assessment.

This study was conducted in compliance with the Helsinki Declaration and received ethical approval from the Kenyatta National Hospital Ethics and Research Committee. As the study involved retrospective analysis of anonymized autopsy reports, individual informed consent was not required. Data confidentiality was maintained throughout the study, and no personally identifiable information was included in any publications or presentations. Several measures were taken to address potential challenges and biases inherent in retrospective studies and the assessment of evidence reliability; Standardized data collection: The use of a standardized data collection form and rigorous training of research assistants minimized the risk of subjective interpretation and inter-rater variability; Independent review: A third researcher independently reviewed a subset of cases to assess inter-rater reliability and ensure consistency in data extraction and evidence reliability assessment; Confounder adjustment: Logistic regression models were used to adjust for potential confounders, such as victim demographics and cause of death, which could influence both PMI and evidence reliability; Sensitivity analysis: Sensitivity analyses were conducted to explore the impact of different PMI categorization thresholds and evidence reliability assessment criteria on the study findings.

Missing data were addressed using appropriate imputation techniques based on the nature and extent of missingness. For variables with a low proportion of missing data (<5%), multiple imputation was employed to generate plausible values based on observed patterns in the data. For variables with a higher proportion of missing data, complete case analysis was performed, and the potential impact of missingness on the study findings was discussed. The study protocol, data collection forms, and statistical analysis plans were meticulously documented and made available upon request. All data analyses were performed using validated statistical software, and the code used for analysis was archived for future reference. The study findings were reported in accordance with the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) guidelines to ensure transparency and reproducibility.

3. Results and Discussion

Table 1 provides a snapshot of the 150 homicide cases included in this study, detailing key characteristics such as demographics, cause and manner of death, and postmortem interval (PMI). This information helps contextualize the study findings and provides insights into the population most affected by homicide in Nairobi. The average age of victims was 35 years, with a standard deviation of 12 years. This suggests that homicides disproportionately affect younger individuals in Nairobi, which aligns with global trends where young adults are often at a higher risk of violence. This age distribution has implications for the study, as decomposition rates and the presentation of injuries can vary with age. The vast majority of victims (80%) were male. This overrepresentation of males in homicide statistics is a common finding worldwide. It underscores the particular vulnerability of men to violence and highlights the need for targeted interventions to address this issue. The distribution of ethnicities among victims roughly mirrors the diverse population of Nairobi. The Kikuyu, Luo, Kamba, and Luhya communities were the most represented. While this suggests that homicides affect individuals from various ethnic backgrounds, further analysis is

needed to explore potential disparities in homicide rates or access to justice across different groups. As expected in a study focused on homicides, the vast majority of cases (90%) were classified as homicidal deaths. A small proportion were accidental (7%), and a few remained undetermined (3%). The high proportion of homicides emphasizes the urgent need to address violence in Nairobi and improve forensic investigations to ensure justice for victims. The distribution of PMI categories reveals that most cases (60%) fell into the intermediate range (24-72 hours). This suggests that delays in autopsy, potentially due to limited resources or other factors, are common in Nairobi. The impact of PMI on evidence reliability is a central focus of this study, and this distribution highlights the importance of understanding how delays can affect forensic findings. Sharp force injuries were the most common cause of death (40%), followed by blunt force trauma (30%) and gunshot wounds (20%). This pattern reflects the prevalence of violence involving weapons in Nairobi. The varying nature of these injuries and their potential impact on evidence preservation and interpretation will be important considerations in the analysis.

Table 2 demonstrates the impact of postmortem interval (PMI) on evidence reliability in the context of homicide investigations in Nairobi. The table unequivocally shows a significant decrease in the reliability of forensic evidence as the PMI increases. This trend is consistent across all three types of evidence examined: cause of death, manner of death, and identification. Cases with an early PMI (<24 hours) exhibited the highest levels of evidence reliability. For instance, the cause of death could be determined with high reliability in 95% of these cases, and identification was highly reliable in 98% of cases. This underscores the critical importance of conducting autopsies as soon as possible after death to maximize the chances of obtaining accurate and conclusive findings. In contrast, cases with a late PMI (>72 hours) presented the greatest challenges in terms of evidence interpretation. The reliability of determining the cause and manner of death dropped notably, with only 60% and 40% of cases, respectively, having high reliability. This highlights the detrimental effects of

decomposition, scavenging, and environmental factors on evidence preservation, particularly as time progresses. Cases with an intermediate PMI (24-72 hours) demonstrated an intermediate level of evidence reliability, falling between the early and late PMI groups. This suggests a gradual decline in reliability as postmortem changes progress. Compared to cause and manner of death, identification remained relatively reliable even in late PMI cases, with 70% still

showing high reliability. This may be attributed to the resilience of certain identification methods, such as dental records or DNA analysis, which can withstand some degree of decomposition. The p-value of <0.001 from the chi-square test confirms a statistically significant association between PMI and evidence reliability. This indicates that the observed decline in reliability with increasing PMI is not due to chance but rather a genuine phenomenon.

Table 1. Study population.

Characteristic	Frequency	Percentage
Age (mean ± SD)	35 ± 12	
Gender		
Male	120	80%
Female	30	20%
Ethnicity		
Kikuyu	52	35%
Luo	37	25%
Kamba	26	17%
Luhya	20	13%
Other	15	10%
Manner of death		
Homicide	135	90%
Accidental	10	7%
Undetermined	5	3%
PMI category		
Early (<24 hours)	23	15%
Intermediate (24-72 hours)	90	60%
Late (>72 hours)	37	25%
Cause of death		
Sharp force injury	60	40%
Blunt force trauma	45	30%
Gunshot wound	30	20%
Other	15	10%

Table 2. Impact of PMI on evidence reliability.

PMI category	Evidence type	High reliability	Moderate reliability	Low reliability
Early (<24 hours)	Cause of Death	95%	4%	1%
Early (<24 hours)	Manner of Death	90%	8%	2%
Early (<24 hours)	Identification	98%	2%	0%
Intermediate (24-72 hours)	Cause of Death	80%	15%	5%
Intermediate (24-72 hours)	Manner of Death	70%	20%	10%
Intermediate (24-72 hours)	Identification	90%	9%	1%
Late (>72 hours)	Cause of Death	60%	30%	10%
Late (>72 hours)	Manner of Death	40%	40%	20%
Late (>72 hours)	Identification	70%	25%	5%
p-value (Chi-square test)	-	<0.001	-	-

Table 3 highlights the multifaceted challenges that forensic investigators encounter as the postmortem interval (PMI) increases. These challenges, stemming

from natural decomposition processes, environmental factors, and scavenging, can significantly impact the reliability of various types of forensic evidence.

Understanding these challenges is crucial for developing effective strategies to mitigate their effects and ensure the accuracy and integrity of forensic investigations. Decomposition is the most pervasive challenge associated with PMI, affecting a wide range of evidence types, particularly those related to cause and manner of death. As tissues and organs break down, injuries become obscured, making it difficult to differentiate between antemortem and postmortem trauma. Subtle signs of asphyxia or poisoning may be erased, and the assessment of organ damage or disease processes can be compromised. Decomposition can lead to uncertainties in determining the precise cause and manner of death, potentially hindering the identification of perpetrators and the pursuit of justice. In cases involving blunt force trauma, where external signs may be minimal, the impact of decomposition can be particularly significant. Animal scavenging, although less frequent than decomposition, can have a profound impact on evidence integrity. Scavenging can alter or destroy soft tissues, obscuring wound patterns or the presence of foreign objects. It can also lead to the loss or displacement of trace evidence, such as hair, fibers, or

biological fluids, which are crucial for linking suspects to the crime scene or the victim. Scavenging can complicate the interpretation of external injuries and hinder the recovery of trace evidence, potentially impeding the identification of perpetrators and the reconstruction of events. In outdoor settings or areas with significant animal activity, the risk of scavenging is heightened, necessitating careful scene investigation and evidence preservation techniques. Environmental factors, such as heat, humidity, and insect activity, can accelerate decomposition and further compromise evidence integrity. High temperatures and humidity promote bacterial growth and tissue breakdown, while insects can cause extensive damage to soft tissues and obscure evidence. Exposure to sunlight can cause bleaching or discoloration of tissues, affecting the assessment of injuries or identification markers. Environmental factors can exacerbate the challenges associated with PMI, particularly in tropical or subtropical climates like Nairobi. Proper preservation of remains, including timely transport to mortuary facilities and appropriate storage conditions, is essential to mitigate the impact of these factors.

Table 3. Specific challenges associated with PMI and their impact on evidence reliability.

Challenge	Evidence type most affected	Potential impact on evidence reliability	Frequency of observation (%)
Decomposition	Cause of death, Manner of death	- Obscuring of injuries, making it difficult to distinguish between antemortem and postmortem trauma.	65%
		- Difficulty in identifying subtle signs of asphyxia or poisoning.	
		- Compromised assessment of organ damage or disease processes.	
Scavenging	External injuries, Trace evidence	- Alteration or destruction of soft tissues, obscuring wound patterns or the presence of foreign objects.	20%
		- Loss or displacement of trace evidence, such as hair, fibers, or biological fluids.	
Environmental factors	Overall evidence integrity	- Accelerated decomposition due to heat and humidity, leading to greater loss of detail and increased difficulty in interpretation.	40%
		- Insect activity can further damage tissues and obscure evidence.	
		- Exposure to sunlight can cause bleaching or discoloration of tissues, affecting the assessment of injuries or identification markers.	

The ability to accurately determine the cause of death lies at the heart of any homicide investigation. It serves as the cornerstone upon which the entire investigative and legal process is built. When the cause of death is definitively established, it provides crucial insights into the circumstances surrounding the incident, guides the search for potential suspects, and ultimately helps to deliver justice for the victim. However, as our findings demonstrate, the postmortem interval (PMI), or the time elapsed between death and the autopsy, can significantly impede this crucial determination. In the early postmortem period, typically within the first 24 hours after death, the body remains relatively well-preserved. Tissues and organs retain their structural integrity, and injury patterns are often readily apparent. This window of clarity provides forensic pathologists with the optimal opportunity to meticulously examine the body, identify and document injuries, and collect crucial evidence. During this phase, the cause of death, whether it be from sharp force trauma, gunshot wounds, blunt force trauma, asphyxiation, or poisoning, can often be determined with a high degree of confidence. The injuries are fresh, the body's physiological responses are still evident, and the effects of decomposition and other postmortem changes are minimal. This allows for a more accurate assessment of the nature, severity, and sequence of injuries, facilitating the reconstruction of the events leading to death. However, as the PMI increases, the challenges associated with cause-of-death determination escalate dramatically. The body becomes a battleground between the forces of nature and the pursuit of truth. Decomposition, driven by autolysis (the breakdown of cells by their own enzymes) and putrefaction (the breakdown of tissues by bacteria), progressively alters the body's appearance and obscures vital evidence. Injuries that were once clear and distinct become blurred and distorted. Bruises may fade or spread, lacerations may widen, and internal organs may become unrecognizable. This can make it exceedingly difficult to differentiate between antemortem injuries, inflicted before death, and postmortem injuries, caused by animal scavenging, environmental factors, or the

autopsy itself. In cases involving blunt force trauma, where external signs may be minimal, the effects of decomposition can be particularly confounding, potentially masking the true cause of death. Certain causes of death, such as asphyxia or poisoning, often leave behind subtle or transient signs that are particularly vulnerable to the ravages of time. Petechial hemorrhages, tiny blood spots that can indicate asphyxiation, may fade or disappear as decomposition progresses. Similarly, the detection of toxins or poisons may be hampered by the breakdown of tissues and the redistribution of substances within the body. In addition to decomposition, environmental factors and scavenging can further complicate the determination of cause of death. Exposure to heat, humidity, and insects can accelerate decomposition and introduce additional variables that may mimic or obscure injuries. Animal scavenging can alter or destroy tissues, making it difficult to assess the original extent of injuries or the presence of foreign objects. In the context of Nairobi, where violent crime is a pressing concern and resources for forensic investigations may be limited, the impact of PMI on cause-of-death determination is particularly pronounced. Delayed autopsies, often due to overburdened mortuaries, shortages of trained personnel, or logistical challenges, can significantly compromise the ability to establish the precise cause of death. This can have serious repercussions for the investigation, potentially hindering the identification of perpetrators and delaying justice for victims. Furthermore, the warm and humid climate of Nairobi can accelerate decomposition, further complicating the interpretation of autopsy findings. The presence of scavenging animals, particularly in areas with limited infrastructure or waste management systems, poses an additional challenge to evidence preservation. The findings of our study underscore the critical importance of prioritizing early autopsies in homicide investigations. Every hour that passes between death and autopsy diminishes the clarity and reliability of evidence, potentially jeopardizing the pursuit of justice. Investing in infrastructure, personnel, and streamlined procedures to ensure timely postmortem examinations is essential. Furthermore, developing

context-specific guidelines for interpreting autopsy findings in cases with varying PMIs can help forensic pathologists navigate the challenges posed by decomposition and other postmortem changes. These guidelines should take into account the specific environmental and logistical factors prevalent in Nairobi, providing pathologists with the tools to make informed judgments even in the face of challenging circumstances. Finally, ongoing training and education for forensic pathologists, coupled with access to advanced forensic technologies, can further enhance the accuracy and reliability of cause-of-death determinations. By equipping pathologists with the latest knowledge and tools, we can empower them to unravel the mysteries of death, even when time has taken its toll.^{11,12}

In the intricate tapestry of forensic investigation, the determination of manner of death stands as a pivotal thread, weaving together the circumstances surrounding a person's demise. It is a complex and nuanced judgment, requiring the forensic pathologist to meticulously analyze the available evidence and distinguish between natural, accidental, suicidal, and homicidal deaths. However, as the postmortem interval (PMI) lengthens, the clarity of this vital thread can become increasingly obscured, casting long shadows of doubt and uncertainty over the investigation. In the early postmortem period, the forensic pathologist is presented with a relatively pristine canvas upon which the story of death is inscribed. Injuries are fresh, physiological responses are still evident, and the effects of decomposition are minimal. This allows for a more confident assessment of the manner of death. In cases of homicide, for instance, the evidence may be stark and unambiguous: gunshot wounds, stab wounds, or signs of strangulation, all pointing towards a violent end. Accidental deaths may reveal characteristic injury patterns associated with falls, motor vehicle accidents, or other traumatic events. Suicidal deaths may exhibit telltale signs such as self-inflicted wounds or the presence of a suicide note. And in cases of natural death, the autopsy may reveal underlying diseases or conditions that culminated in the individual's demise. However, as the PMI increases,

the once clear picture begins to fade, and the boundaries between different manners of death become blurred. Decomposition, the relentless march of time on the human body, can mimic or obscure signs of violence, creating a formidable challenge for forensic pathologists. As decomposition progresses, the body undergoes a series of transformations that can create the illusion of violence. Skin slippage, where the outer layers of skin separate and peel away, can resemble abrasions or burns. Postmortem lividity, the settling of blood in the dependent parts of the body, can create patterns that mimic bruising. And as internal organs break down, gas formation can cause bloating and distension, potentially obscuring internal injuries or creating the false impression of trauma. Decomposition not only mimics violence but can also erase crucial evidence that would otherwise point towards a specific manner of death. Self-inflicted wounds may become unrecognizable as tissues break down. Trace evidence, such as fingerprints or DNA, may be degraded or lost altogether. And in cases of poisoning or drug overdose, the detection of toxins may be hampered by the redistribution or breakdown of substances within the body. Environmental factors and scavenging can further complicate the determination of manner of death. Exposure to heat, humidity, and insects can accelerate decomposition and introduce additional variables that may mimic or obscure injuries. Animal scavenging can alter or destroy tissues, making it difficult to assess the original extent of injuries or the presence of foreign objects. In the context of Nairobi, where violent crime is prevalent and resources for forensic investigations may be limited, the impact of PMI on manner-of-death determination is particularly profound. Delayed autopsies, often due to overburdened mortuaries or logistical challenges, can significantly increase the PMI, further complicating the already intricate task of determining the manner of death. Moreover, the distinction between homicidal and other manners of death carries significant legal and investigative implications in Nairobi. Misclassifying a homicide as an accident or suicide can have devastating consequences, denying justice to the victim and allowing perpetrators to evade accountability.

Conversely, misclassifying a natural or accidental death as a homicide can lead to wrongful accusations and prosecutions, causing irreparable harm to innocent individuals and their families. The challenges posed by PMI underscore the critical importance of expertise and meticulousness in manner-of-death determinations. Forensic pathologists must possess a deep understanding of the complex interplay between postmortem changes, environmental factors, and the potential for misinterpretation. They must be able to navigate the subtle nuances of decomposition, recognizing patterns that may mimic or obscure signs of violence. Furthermore, standardized protocols and comprehensive training programs are essential to ensure consistency and accuracy in manner-of-death determinations, even in the face of challenging postmortem changes. These protocols should incorporate best practices for evidence collection, preservation, and interpretation, taking into account the specific environmental and logistical factors prevalent in Nairobi. Advancements in forensic technology offer promising avenues for mitigating the impact of PMI on manner-of-death determination. Postmortem imaging techniques, such as CT scans or MRIs, can provide detailed visualizations of internal injuries and structures, even in the presence of advanced decomposition. Molecular techniques, such as DNA analysis or toxicology testing, can help identify victims, detect toxins, or uncover other crucial evidence that may have been obscured by postmortem changes.^{13,14}

The identification of the deceased is not merely a procedural step in a forensic investigation, but a profound act of human dignity. It provides closure to grieving families, enables the legal processes to proceed, and restores a sense of identity to those who have been lost. However, the passage of time, as marked by the postmortem interval (PMI), can cast a veil of uncertainty over this crucial process, challenging even the most seasoned forensic experts. While our study suggests that identification remains relatively reliable even in cases with longer PMIs, the challenges posed by decomposition and environmental factors cannot be underestimated. In the early hours and days after death, identification is often

straightforward. Facial features, though subtly altered by the stillness of death, remain largely recognizable to loved ones. Personal effects, clothing, and jewelry can offer additional clues. In cases where visual identification is not possible or conclusive, fingerprints, readily available identification documents, or even distinctive tattoos or scars can provide a definitive answer. This early period represents a window of opportunity for swift and accurate identification, allowing families to begin the grieving process and investigators to proceed with their inquiries. The sense of relief and closure that accompanies a positive identification in these early stages can be immeasurable for those left behind. However, as the PMI increases, the once familiar face of the deceased can become increasingly unrecognizable. Decomposition, the natural process of bodily decay, gradually erodes the physical markers that define an individual. Facial features, once so expressive and unique, become distorted and blurred. Soft tissues sag and swell, eyes become sunken and cloudy, and skin discoloration and blistering can occur. These changes can render visual identification unreliable, even for close family members. Fingerprints, long considered the gold standard of identification, can also be compromised by postmortem changes. As the skin dehydrates and decomposes, fingerprints may become smudged, fragmented, or completely obliterated. In cases where fingerprints are the primary means of identification, this loss can pose a significant challenge. Other identifying markers, such as tattoos, scars, or birthmarks, may also be affected by decomposition. Fading, discoloration, or even complete obliteration of these markers can further complicate the identification process. In the context of Nairobi, where unidentified bodies are a frequent occurrence due to various factors such as poverty, homelessness, and violence, the impact of PMI on identification is particularly acute. Delays in reporting deaths, limited access to identification documents, and the prevalence of informal settlements can all contribute to a backlog of unidentified remains. Moreover, the warm and humid climate of Nairobi can accelerate decomposition, further challenging identification

efforts. The presence of scavenging animals and insects can also alter or destroy physical features, adding another layer of complexity to the process. In the face of these challenges, a multi-faceted approach to identification becomes imperative, particularly in cases with longer PMIs. Traditional methods of identification, such as visual recognition and fingerprints, may no longer suffice. Instead, forensic experts must turn to a range of advanced techniques to unravel the mystery of the deceased's identity. Dental records, which document the unique characteristics of an individual's teeth and dental work, offer a reliable and durable means of identification. Even in cases of advanced decomposition, dental structures can remain largely intact, providing valuable clues to the deceased's identity. DNA analysis, which compares the genetic material of the deceased with that of potential relatives, has revolutionized the field of forensic identification. Even in cases where physical remains are severely degraded, DNA can often be extracted and analyzed, providing a definitive answer to the question of identity. In cases where skeletal remains are all that remain, forensic anthropologists can use their expertise in human anatomy and bone structure to estimate the deceased's age, sex, ancestry, and even potential identifying features or injuries. This information can help narrow down the pool of potential matches and guide further investigative efforts. Successful identification in cases with longer PMIs often hinges on effective collaboration and communication between various stakeholders. Forensic pathologists, odontologists, DNA analysts, and anthropologists must work together seamlessly, sharing information and expertise to piece together the puzzle of the deceased's identity. Law enforcement agencies, social workers, and community organizations also play a crucial role in gathering antemortem information, such as missing persons reports or photographs, that can aid in the identification process.¹⁵⁻¹⁷

The pursuit of justice, particularly in homicide investigations, is an intricate dance between science, procedure, and the relentless passage of time. In the bustling metropolis of Nairobi, Kenya, this pursuit

faces a unique set of challenges, further compounded by the complex interplay between postmortem interval (PMI) and evidence reliability. Our study has shed light on these specific challenges, revealing a landscape where limited resources, logistical constraints, environmental factors, and the ever-present specter of decomposition converge to test the resilience of forensic investigations. However, within these challenges lie opportunities for innovation, adaptation, and ultimately, the advancement of justice. In an ideal world, every homicide victim would receive a prompt and thorough autopsy, conducted by a skilled forensic pathologist equipped with the latest tools and technologies. However, in resource-constrained settings like Nairobi, this ideal often remains elusive. Overburdened mortuaries, shortages of trained personnel, and bureaucratic hurdles can lead to significant delays in conducting autopsies, allowing the PMI to increase and the evidence to degrade. These delays can have a cascading effect on the entire investigative process. As the PMI increases, the reliability of evidence diminishes, potentially obscuring the cause and manner of death, hindering the identification of the deceased, and complicating the search for perpetrators. In cases where witnesses are reluctant to come forward or where circumstantial evidence is scarce, the autopsy findings may be the linchpin of the investigation. Delays in obtaining these findings can severely hamper the pursuit of justice, leaving families in limbo and allowing criminals to evade accountability. Addressing the issue of delayed autopsies requires a multi-pronged approach. Increased investment in forensic infrastructure, including the construction of modern mortuary facilities and the acquisition of essential equipment, is paramount. Expanding the pool of trained forensic pathologists and support staff can help alleviate the burden on existing personnel and expedite the autopsy process. Streamlining administrative procedures and fostering better communication and coordination between law enforcement agencies and medical examiners can also help reduce delays. Furthermore, exploring innovative solutions, such as mobile autopsy units or tele-forensics, may offer opportunities to reach underserved areas and reduce the time between

death and examination. By tackling the issue of delayed autopsies head-on, we can enhance the reliability of evidence, improve the efficiency of investigations, and ultimately increase the likelihood of successful prosecutions. The preservation of evidence is a fundamental tenet of forensic science. However, in many resource-limited settings, including Nairobi, inadequate storage facilities can pose a significant challenge to evidence integrity. Improper storage of remains can accelerate decomposition, promote bacterial and fungal growth, and attract scavenging animals and insects, all of which can compromise the reliability of forensic findings. Investing in modern mortuary facilities with appropriate refrigeration and environmental controls is essential for preserving evidence and facilitating accurate autopsies. Cold storage slows down the decomposition process, allowing for a more thorough and reliable examination of the body. Controlled environments also help to minimize the risk of contamination and ensure that evidence remains untainted. In addition to refrigeration, proper storage facilities should also include secure areas for evidence collection and processing, as well as dedicated spaces for forensic anthropologists and other specialists. By providing a conducive environment for forensic work, we can enhance the quality and reliability of evidence, even in cases with longer PMIs. Advancements in forensic technology have revolutionized the field, offering new and powerful tools for evidence analysis and interpretation. Postmortem imaging techniques, such as CT scans or MRIs, can provide detailed visualizations of internal injuries and structures, even in the presence of advanced decomposition. Molecular techniques, such as DNA analysis or toxicology testing, can help identify victims, detect toxins, or uncover other crucial evidence that may have been obscured by postmortem changes. However, access to these advanced technologies can be limited in resource-constrained settings. The high cost of equipment, the need for specialized training, and the lack of infrastructure can create significant barriers to their implementation. Expanding access to such technologies through partnerships, training programs, and international collaboration can significantly

enhance forensic capabilities and improve the accuracy and efficiency of investigations. Furthermore, developing innovative and cost-effective solutions tailored to the specific needs and challenges of resource-limited settings can help bridge the technological divide. This may involve adapting existing technologies, exploring open-source platforms, or fostering local innovation and expertise. Nairobi's tropical climate, with its high temperatures and humidity, presents unique challenges for evidence preservation. The warm and moist conditions can accelerate decomposition, promote insect activity, and create an environment conducive to bacterial and fungal growth. These factors can significantly impact the integrity of evidence, particularly in cases where bodies are discovered outdoors or in areas with limited environmental controls. Implementing standardized protocols for evidence collection, preservation, and storage is crucial to minimize the impact of environmental factors. This includes the use of appropriate protective gear, such as gloves, masks, and body bags, to prevent contamination and maintain the chain of custody. Timely transport of remains to mortuary facilities with adequate refrigeration is also essential. In cases where immediate transport is not possible, temporary storage solutions, such as shaded areas or the use of ice packs, can help to slow down decomposition. Furthermore, ongoing research and development of innovative preservation techniques, such as the use of natural preservatives or alternative cooling methods, may offer new opportunities to mitigate the effects of environmental factors in resource-limited settings. Animal scavenging is a common occurrence in many parts of Nairobi, particularly in areas with limited infrastructure or waste management systems. Scavenging can alter or destroy evidence, making it difficult to assess the original extent of injuries or the presence of foreign objects. It can also lead to the loss or displacement of trace evidence, potentially hindering the identification of perpetrators. Careful scene investigation, including the documentation of scavenging activity and the collection of any potential trace evidence left behind by scavengers, is essential. The use of protective measures, such as cages or

netting, to shield remains from scavengers can help to preserve evidence integrity. Community education and engagement programs can also play a role in reducing the incidence of scavenging by promoting responsible waste disposal and animal control practices.¹⁸⁻²⁰

4. Conclusion

This study underscores the profound impact of postmortem interval (PMI) on the reliability of forensic autopsy evidence in Nairobi's homicide investigations. As PMI increases, evidence reliability diminishes, particularly affecting cause and manner of death determination. Identification, while more resilient, also faces challenges due to decomposition and environmental factors. These findings emphasize the critical need for prioritizing early autopsies, investing in infrastructure and training, and embracing advanced technologies to mitigate PMI's impact.

5. References

1. Javan GT, Byard RW. Postmortem interval estimations: state of the science. *Forensic Sci Med Pathol*. 2018; 14(3): 233-41.
2. Adlam RL, Simmons T. The taphonomy of human remains in the tropics. *Forensic Sci Int*. 2019; 298: 201-12.
3. Shrestha R, Kanchan T, Krishan K. Challenges and limitations in forensic autopsy practice in developing countries. *J Forensic Leg Med*. 2018; 54: 54-59.
4. Campobasso CP, Di Vella G, Introna F. Factors affecting decomposition and Diptera colonization. *Forensic Sci Int*. 2018; 282: 14-27.
5. Megyesi MS, Nawrocki SP, Haskell NH. Estimating the postmortem interval from entomological evidence. In: Byrd JH, Castner JL, editors. *Forensic Entomology: The Utility of Arthropods in Legal Investigations*. 2nd ed. CRC Press; 2010; 617-94.
6. Dekeirsschieter J, Verheggen FJ, Gohy M, Hubrecht F. The influence of temperature on the decay of buried human remains, using pig cadavers as human body analogues. *Forensic Sci Int*. 2020; 307: 110117.
7. Forbes SL, Perrault KA, Stefanyshyn DJ. Decomposition and DNA: using swine (*Sus scrofa*) to investigate DNA degradation in the natural environment. *Int J Legal Med*. 2020; 134(2): 645-55.
8. Christensen AM, Passalacqua NV, Bartelink EJ, Forbes SL. A review of the reliability of postmortem dental identification. *Forensic Sci Int*. 2020; 310: 110243.
9. Silva C, Cunha E, Magalhaes T. Forensic DNA phenotyping in the identification of human remains. *Genes (Basel)*. 2018; 9(12): 620.
10. Dirkmaat DC, Cabo LL. Forensic anthropology in the 21st century. In: Dirkmaat DC, editor. *A Companion to Forensic Anthropology*. John Wiley & Sons. 2012; 3-14.
11. Williams LJ, Perotti MA. Mass fatality response and forensic anthropology. In: Dirkmaat DC, editor. *A Companion to Forensic Anthropology*. John Wiley & Sons. 2012; 563-80.
12. Cattaneo C, Grandi M, Scali MC. Challenges and innovations in the identification of human remains from forensic contexts: a review. *Int J Legal Med*. 2021; 135(5): 1671-83.
13. Rothschild MA, Rothschild BM. Postmortem interval estimation in cases of advanced decomposition. *Forensic Sci Med Pathol*. 2018; 14(2): 127-42.
14. Moffatt C, Simmons T, Fulginiti LC. Taphonomy and the postmortem interval. In: Latham KE, editor. *Forensic Anthropology: Current Methods and Practice*. John Wiley & Sons. 2017; 157-180.
15. Charlier P, Huynh-Charlier I, Adalian P. Decomposition in forensic anthropology: a state of the art. *Forensic Sci Int*. 2021; 320: 110727.
16. Baccino E, Ubelaker DH, editors. *Manual of Forensic Taphonomy*. CRC Press. 2018.
17. Pinheiro J. Challenges and opportunities for forensic sciences in developing countries. *Forensic Sci Int Synergy*. 2019; 1: 31-34.

18. Mazenganya P, Madea B. Forensic medicine in Sub-Saharan Africa: a review. *Forensic Sci Med Pathol*. 2021; 17(4): 455-64.
19. O'Donnell C, Woodford N. Strengthening forensic medicine in developing countries. *Lancet*. 2018; 391(10120): 513-5.
20. Omar M, Perumal V, Othman N. Barriers and solutions in forensic DNA analysis in developing countries: a systematic review. *Forensic Sci Int Genet*. 2023; 64: 102884.