



## Exploring the Role of Inflammatory Cytokines and Oxidative Stress in Febrile Seizure Pathogenesis: A Case-Control Study in Mumbai, India

Lakshmi Srivanathan<sup>1\*</sup>, Saeed Khan<sup>1</sup>

<sup>1</sup>Department of Pediatric Neurology, Mumbai Hospital, India

### ARTICLE INFO

#### Keywords:

Children  
Febrile seizures  
Inflammatory cytokines  
India  
Oxidative stress

#### \*Corresponding author:

Lakshmi Srivanathan

#### E-mail address:

[lakshmisr@gmail.com](mailto:lakshmisr@gmail.com)

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.59345/sjped.v1i2.66>

### ABSTRACT

**Introduction:** Febrile seizures (FS) are the most common childhood seizure disorder, often causing parental anxiety and posing a challenge for healthcare professionals. While the exact pathogenesis remains unclear, recent research suggests a complex interplay of genetic predisposition, fever, and inflammatory processes. This study aimed to investigate the role of inflammatory cytokines and oxidative stress markers in FS pathogenesis among children in Mumbai, India. **Methods:** A case-control study was conducted involving 100 children with FS (cases) and 100 age-matched febrile children without seizures (controls) admitted to a tertiary care hospital in Mumbai. Serum levels of inflammatory cytokines (IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and IL-10) and oxidative stress markers (malondialdehyde [MDA] and superoxide dismutase [SOD]) were measured and compared between the two groups. **Results:** Children with FS exhibited significantly higher serum levels of pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF- $\alpha$  ( $p < 0.001$  for all) compared to controls. Conversely, the anti-inflammatory cytokine IL-10 was significantly lower in the FS group ( $p < 0.001$ ). Furthermore, MDA levels were significantly elevated ( $p < 0.001$ ), while SOD levels were significantly decreased ( $p < 0.001$ ) in the FS group compared to controls. **Conclusion:** This study provides evidence for the involvement of inflammatory cytokines and oxidative stress in FS pathogenesis. Elevated pro-inflammatory cytokines and oxidative stress markers, coupled with decreased anti-inflammatory cytokine levels, suggest a dysregulated inflammatory response and impaired antioxidant defense mechanism in children with FS.

### 1. Introduction

Febrile seizures (FS), the most prevalent neurological disorder in young children, represent a significant concern for both families and healthcare providers worldwide. Affecting children typically between 6 months and 5 years of age, FS is characterized by convulsions triggered by a febrile illness, in the absence of central nervous system infection or any other identifiable cause. While generally considered benign and self-limiting, the experience can be profoundly distressing for parents and caregivers. Moreover, a small subset of children with FS may face an increased risk of developing epilepsy later in life. Understanding the pathogenesis

of FS is crucial for developing effective preventive and therapeutic strategies. However, despite extensive research, the exact mechanisms underlying FS remain elusive. Current understanding suggests a complex interplay of factors, including genetic predisposition, fever, and inflammatory processes. This intricate interplay necessitates a multifaceted approach to research, encompassing genetic, physiological, and immunological perspectives.<sup>1,2</sup>

Genetic factors are believed to play a significant role in FS susceptibility. Numerous studies have identified genes involved in ion channel function and neuronal excitability that are implicated in FS. These genes, often associated with familial forms of epilepsy,

highlight the inherent vulnerability of certain individuals to seizures in the context of fever. Mutations in genes encoding voltage-gated sodium channels (e.g., SCN1A, SCN1B) and GABA receptors have been particularly associated with FS and epilepsy syndromes. Such genetic variations can alter neuronal excitability, making individuals more susceptible to seizures when challenged by fever. Fever, often caused by viral or bacterial infections, is the primary trigger for FS. The rapid rise in body temperature associated with fever is thought to alter neuronal excitability, leading to seizures. This alteration in neuronal function may be mediated by changes in ion channel kinetics, neurotransmitter release, and synaptic plasticity. Furthermore, fever can induce the production of inflammatory mediators, further contributing to the pathogenesis of FS.<sup>3,4</sup>

In recent years, increasing evidence has pointed towards the involvement of inflammatory processes in FS pathogenesis. Studies have demonstrated elevated levels of pro-inflammatory cytokines, such as interleukin-1 $\beta$  (IL-1 $\beta$ ), interleukin-6 (IL-6), and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), in the serum and cerebrospinal fluid of children with FS. These cytokines are key mediators of the inflammatory response and can affect neuronal excitability by modulating neurotransmitter release and ion channel function. IL-1 $\beta$ , a potent pro-inflammatory cytokine, is produced by activated immune cells in response to infection or injury. It can increase neuronal excitability by promoting glutamate release, the major excitatory neurotransmitter in the brain, and inhibiting GABAergic neurotransmission, the primary inhibitory neurotransmitter system. IL-6, another pro-inflammatory cytokine, can induce neuronal hyperexcitability and contribute to seizure activity. TNF- $\alpha$ , a pleiotropic cytokine with both pro- and anti-inflammatory properties, can also enhance neuronal excitability and promote inflammation in the brain. The elevated levels of pro-inflammatory cytokines observed in children with FS suggest a dysregulated inflammatory response that may contribute to neuronal hyperexcitability and lower the seizure threshold. This dysregulation may be driven by the underlying infection or fever, or it may reflect an

inherent predisposition to exaggerated inflammatory responses in some individuals.<sup>5,6</sup>

In addition to inflammation, oxidative stress has also been implicated in FS pathogenesis. Oxidative stress refers to an imbalance between the production of reactive oxygen species (ROS) and the body's antioxidant defense mechanisms. Excessive ROS production can lead to cellular damage and dysfunction, including neuronal injury, which may contribute to seizure activity. ROS are highly reactive molecules that can damage cellular components, such as lipids, proteins, and DNA, leading to neuronal dysfunction and death. Under normal physiological conditions, ROS are produced as byproducts of cellular metabolism and play important roles in various signaling pathways. However, during infection or inflammation, ROS production can increase dramatically, overwhelming the body's antioxidant defenses. Antioxidant enzymes, such as superoxide dismutase (SOD), catalase, and glutathione peroxidase, play a crucial role in neutralizing ROS and protecting cells from oxidative damage. SOD, a key antioxidant enzyme, catalyzes the dismutation of superoxide radicals into hydrogen peroxide, which is then further detoxified by catalase and glutathione peroxidase. Studies have reported increased levels of oxidative stress markers, such as malondialdehyde (MDA), a marker of lipid peroxidation, and decreased levels of antioxidant enzymes, such as SOD, in children with FS. These findings suggest that oxidative stress may contribute to neuronal injury and seizure activity in FS.<sup>7,8</sup> While numerous studies have investigated the role of inflammatory cytokines and oxidative stress in FS, most of this research has been conducted in developed countries. Limited data is available from developing nations like India, where the prevalence of FS is high and healthcare access may be limited. Understanding the specific factors contributing to FS pathogenesis in this context is crucial for developing targeted interventions and improving outcomes for children in India.<sup>9,10</sup> This study aimed to address this gap by investigating the role of inflammatory cytokines and oxidative stress markers in FS pathogenesis among children in Mumbai, India.

## 2. Methods

A hospital-based case-control study was conducted at a tertiary care hospital in Mumbai, India, renowned for its specialized pediatric neurology department and comprehensive diagnostic facilities. The study spanned a year, from January 2023 to December 2023, to capture a representative sample of patients across different seasons and potential variations in infectious disease prevalence. This timeframe allowed for the recruitment of an adequate number of participants while minimizing potential biases associated with seasonal variations in FS incidence. The study population comprised two distinct groups; Cases: 100 children aged between 6 months and 5 years diagnosed with FS; Controls: 100 age-matched febrile children without seizures. The inclusion criteria for cases were meticulously defined to ensure accurate representation of FS; Age: 6 months to 5 years, the age range during which FS are most prevalent; Febrile Seizure Episode: A documented episode of seizure activity concurrent with a febrile illness (temperature  $\geq 38^{\circ}\text{C}$ ), ascertained through detailed parental reporting and medical record review; Absence of CNS Infection: Exclusion of any underlying central nervous system infection, such as meningitis or encephalitis, confirmed by clinical examination, lumbar puncture, and neuroimaging studies when indicated; No Identifiable Cause: Exclusion of other identifiable causes of seizures, including metabolic disorders, electrolyte imbalances, and prior history of afebrile seizures. The control group, crucial for comparison, consisted of children admitted to the pediatric department with fever but without any history of seizures. Age-matching with the case group was meticulously performed to minimize the influence of age-related variations in immune function and oxidative stress parameters. Exclusion criteria, applied rigorously to both groups, aimed to eliminate potential confounding factors; History of Afebrile Seizures: To isolate the specific impact of febrile illness on seizure activity; Presence of CNS Infection: To exclude seizures directly attributable to infectious processes within the central nervous system; Metabolic Disorders: To avoid inclusion of children with underlying metabolic conditions that could

independently influence seizure susceptibility or inflammatory/oxidative stress profiles; Electrolyte Imbalances: To eliminate seizures triggered by electrolyte disturbances, which can mimic FS; Chronic Illnesses: To exclude children with chronic medical conditions that could affect immune function or oxidative stress parameters; Prior Antipyretic/Anticonvulsant Use: To avoid potential masking of inflammatory responses or alteration of seizure thresholds by prior medication use. Recruitment was conducted through the pediatric emergency department and inpatient wards. Potential participants were identified by trained research nurses who screened admissions based on the predefined inclusion and exclusion criteria. Parents or legal guardians of eligible children were approached and provided with detailed information about the study objectives, procedures, and potential risks and benefits. Written informed consent was obtained from parents or legal guardians prior to enrollment.

The study protocol was meticulously reviewed and approved by the Institutional Ethics Committee of the hospital, ensuring adherence to all ethical principles for research involving human subjects. The study was conducted in accordance with the Declaration of Helsinki and national ethical guidelines. Informed consent procedures were designed to be clear, comprehensive, and culturally sensitive, ensuring that parents or guardians fully understood the implications of participation. Confidentiality of participant data was maintained throughout the study, and all data were anonymized to protect individual privacy. A standardized data collection form was developed to ensure consistency and accuracy in data recording. Trained research nurses collected the following information; Demographic Data: Age, gender, socioeconomic status (assessed using a validated socioeconomic scale relevant to the Indian context), and relevant medical history; Clinical Data: Details of the febrile illness, including onset, duration, maximum temperature recorded, and suspected etiology (viral or bacterial). For cases, information on seizure characteristics (type, duration, frequency) was also meticulously documented.

Blood samples were collected from all participants within 24 hours of hospital admission to capture the acute phase of the febrile illness and potential seizure activity. A standardized phlebotomy procedure was followed to minimize discomfort and ensure sample quality. Approximately 5 mL of venous blood was collected into sterile serum separator tubes. The tubes were gently inverted to promote clotting and then centrifuged at 3000 rpm for 10 minutes to separate the serum. Serum samples were carefully aliquoted and stored at -80°C until analysis, ensuring preservation of cytokine and oxidative stress marker integrity.

Serum levels of inflammatory cytokines (IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and IL-10) were quantified using commercially available enzyme-linked immunosorbent assay (ELISA) kits (R&D Systems, USA). These kits are widely recognized for their high sensitivity and specificity in cytokine measurement. All assays were performed in duplicate according to the manufacturer's instructions. Strict quality control measures were implemented, including the use of standard curves and internal controls, to ensure assay accuracy and reliability. Malondialdehyde (MDA), a marker of lipid peroxidation, was measured using a spectrophotometric method based on the reaction with thiobarbituric acid (TBA). This method is well-established and widely used in oxidative stress research. Serum samples were reacted with TBA under acidic conditions, and the resulting MDA-TBA adduct was quantified spectrophotometrically at 532 nm. Superoxide Dismutase (SOD) activity was measured using a spectrophotometric method based on the inhibition of superoxide radical-induced reduction of nitroblue tetrazolium (NBT). This method is widely used to assess SOD activity in biological samples. Serum samples were added to a reaction mixture containing NBT and an enzymatic system generating superoxide radicals. SOD activity was determined by measuring the inhibition of NBT reduction at 560 nm. All laboratory analyses were performed by trained technicians blinded to the participant group assignment (case or control) to minimize potential bias in data interpretation.

Data were analyzed using SPSS software (version 25.0), a comprehensive statistical package widely used

in medical research. Descriptive statistics were used to summarize demographic and clinical characteristics of the study population. Continuous variables were expressed as mean  $\pm$  standard deviation (SD) or median (interquartile range [IQR]), depending on the distribution assessed using histograms and normality tests. Categorical variables were expressed as frequencies and percentages. Group comparisons were performed using appropriate statistical tests; Continuous Variables: Independent samples t-test was used for normally distributed data, while the Mann-Whitney U test was used for non-normally distributed data; Categorical Variables: Chi-square test or Fisher's exact test was used to compare categorical variables between the two groups. Statistical significance was set at a p-value of <0.05. All statistical analyses were conducted by a biostatistician experienced in pediatric research, ensuring rigor and accuracy in data interpretation.

Stringent quality control measures were implemented throughout the study to ensure data integrity and reliability; Training and Standardization: Research nurses and laboratory technicians underwent comprehensive training on study procedures and data collection methods to ensure consistency and minimize inter-observer variability; Data Validation: Data entry was double-checked by independent personnel to minimize errors; Laboratory Quality Control: Internal controls and standard curves were used in all laboratory assays to ensure accuracy and precision. Regular equipment calibration and maintenance were performed to ensure reliable performance.

### **3. Results and Discussion**

Table 1 presents the demographic characteristics of the 100 children with febrile seizures (FS) and the 100 febrile controls enrolled in the study. It provides a clear picture of the study population, allowing us to assess the comparability of the two groups and identify any potential confounding factors. The mean age of children in both the FS group (24.5  $\pm$  12.3 months) and the control group (23.8  $\pm$  11.8 months) is very similar. The p-value of 0.612 indicates that this difference is not statistically significant. This is crucial

because age can influence both immune responses and susceptibility to seizures. The careful age-matching between the groups ensures that any observed differences in inflammatory markers or oxidative stress are not simply due to age discrepancies. The gender distribution is also comparable between the two groups, with slightly more males in the FS group (60%) and slightly more females in the control group (55%). However, this

difference is not statistically significant ( $p=0.485$ ), suggesting that gender is unlikely to be a major confounding factor in this study. The distribution of socioeconomic status across low, middle, and high categories is almost identical between the FS and control groups. The non-significant  $p$ -value (0.678) indicates that socioeconomic factors are unlikely to influence the relationship between inflammatory cytokines, oxidative stress, and FS in this study.

Table 1. Demographic characteristics.

Characteristic	FS Group (n=100)	Control Group (n=100)	p-value
Age (months)	24.5 ± 12.3	23.8 ± 11.8	0.612
Gender (male/female)	60/40	55/45	0.485
Socioeconomic status (low/middle/high)	30/50/20	35/45/20	0.678

Table 2 presents the serum levels of key inflammatory cytokines in children with febrile seizures (FS) and febrile controls. This data provides crucial insights into the inflammatory response associated with FS. Children with FS exhibit significantly higher levels of all three pro-inflammatory cytokines measured; IL-1 $\beta$ : 15.2 ± 6.8 pg/mL in the FS group compared to 8.5 ± 4.2 pg/mL in the control group ( $p<0.001$ ); IL-6: 32.5 ± 10.5 pg/mL in the FS group compared to 18.3 ± 7.9 pg/mL in the control group ( $p<0.001$ ); TNF- $\alpha$ : 28.7 ± 9.2 pg/mL in the FS group compared to 15.6 ± 6.5 pg/mL in the control group ( $p<0.001$ ). These marked elevations in IL-1 $\beta$ , IL-6, and TNF- $\alpha$  strongly suggest a heightened

inflammatory state in children with FS. These cytokines are known to play pivotal roles in the inflammatory cascade, promoting immune cell activation, and fever generation, and potentially influencing neuronal excitability. Conversely, children with FS show significantly lower levels of the anti-inflammatory cytokine IL-10: 12.4 ± 5.3 pg/mL in the FS group compared to 18.6 ± 7.1 pg/mL in the control group ( $p<0.001$ ). This finding suggests a deficiency in the counter-regulatory mechanisms that normally dampen inflammation. IL-10 typically helps to resolve inflammation and limit excessive immune activation. Its reduced levels in the FS group may contribute to the unchecked pro-inflammatory state.

Table 2. Inflammatory cytokines.

Cytokine	FS Group (pg/mL)	Control Group (pg/mL)	p-value
IL-1 $\beta$	15.2 ± 6.8	8.5 ± 4.2	<0.001
IL-6	32.5 ± 10.5	18.3 ± 7.9	<0.001
TNF- $\alpha$	28.7 ± 9.2	15.6 ± 6.5	<0.001
IL-10	12.4 ± 5.3	18.6 ± 7.1	<0.001

Table 3 presents the levels of two key oxidative stress markers, malondialdehyde (MDA) and superoxide dismutase (SOD), in children with febrile seizures (FS) and febrile controls. This data sheds light on the role of oxidative stress in the pathogenesis of

FS. Children with FS have significantly higher levels of MDA (4.8 ± 1.2 nmol/mL) compared to febrile controls (3.2 ± 0.8 nmol/mL) ( $p<0.001$ ). MDA is a byproduct of lipid peroxidation, a process where reactive oxygen species (ROS) damage cell membranes. Elevated MDA

levels indicate increased oxidative damage to lipids, suggesting a heightened state of oxidative stress in children with FS. Conversely, children with FS show significantly lower levels of SOD ( $85.3 \pm 15.6$  U/mL) compared to febrile controls ( $112.5 \pm 20.3$  U/mL)

( $p < 0.001$ ). SOD is an important antioxidant enzyme that neutralizes superoxide radicals, a type of ROS. Reduced SOD activity indicates a diminished capacity to scavenge these harmful free radicals, further contributing to the oxidative stress imbalance.

Table 3. Oxidative stress markers.

Marker	FS Group	Control Group	p-value
MDA (nmol/mL)	$4.8 \pm 1.2$	$3.2 \pm 0.8$	<0.001
SOD (U/mL)	$85.3 \pm 15.6$	$112.5 \pm 20.3$	<0.001

Our study unequivocally demonstrates a significant elevation in serum levels of pro-inflammatory cytokines IL-1 $\beta$ , IL-6, and TNF- $\alpha$  in children with FS. This observation aligns with a growing body of evidence implicating inflammation as a key player in FS pathogenesis. These cytokines, central to the inflammatory cascade, are not mere bystanders in the febrile response, they actively modulate neuronal excitability and influence seizure susceptibility. To fully appreciate the significance of these findings, it's crucial to delve deeper into the multifaceted roles of these cytokines and explore the potential mechanisms underlying their contribution to FS. Interleukin-1 $\beta$  (IL-1 $\beta$ ) is a potent pyrogen and a central orchestrator of the inflammatory response. Produced primarily by activated macrophages and monocytes, IL-1 $\beta$  exerts its effects by binding to the IL-1 receptor, triggering a cascade of intracellular signaling events that culminate in the production of other inflammatory mediators, such as prostaglandins and chemokines. These mediators contribute to the cardinal signs of inflammation: redness, swelling, heat, and pain. In the context of FS, IL-1 $\beta$ 's role extends beyond its pyrogenic and inflammatory effects. It has been shown to directly influence neuronal excitability, making neurons more susceptible to firing and lowering the seizure threshold. IL-1 $\beta$  can increase the release of glutamate, the primary excitatory neurotransmitter in the brain. Glutamate binds to its receptors on neurons, triggering depolarization and increasing the likelihood of action potential generation. By enhancing glutamate release, IL-1 $\beta$  promotes neuronal excitation

and increases seizure susceptibility. GABA, the main inhibitory neurotransmitter in the brain, counteracts glutamate's excitatory effects by hyperpolarizing neurons and reducing their firing rate. IL-1 $\beta$  can disrupt GABAergic neurotransmission by reducing GABA synthesis, inhibiting GABA receptor function, or promoting GABA reuptake. This disruption of inhibitory signaling further tilts the balance towards excitation, contributing to seizure activity. IL-1 $\beta$  can also directly modulate the activity of ion channels, which are crucial for regulating neuronal excitability. For instance, it can enhance the function of voltage-gated sodium channels, which are responsible for the rapid depolarization phase of action potentials. This increased sodium channel activity can lead to hyperexcitability and increased seizure susceptibility. Astrocytes, the most abundant glial cells in the brain, play a crucial role in maintaining neuronal homeostasis and regulating synaptic transmission. IL-1 $\beta$  can activate astrocytes, leading to the release of pro-inflammatory mediators and further contributing to neuronal dysfunction and seizure activity. Interleukin-6 (IL-6) is another prominent pro-inflammatory cytokine that plays a multifaceted role in FS pathogenesis. Produced by a variety of cells, including immune cells, endothelial cells, and fibroblasts, IL-6 exerts its effects by binding to its receptor, activating intracellular signaling pathways that regulate gene expression and cellular function. Like IL-1 $\beta$ , IL-6 exhibits pro-convulsant properties, contributing to neuronal hyperexcitability and seizure generation. IL-6 can alter the function of various ion channels, including voltage-gated sodium channels,

potassium channels, and calcium channels. These alterations can disrupt the delicate balance of ionic currents that regulate neuronal excitability, leading to hyperexcitability and increased seizure susceptibility. IL-6 can influence the synthesis, release, and reuptake of neurotransmitters, potentially disrupting the delicate balance between excitation and inhibition in the brain. For example, it can increase glutamate release and inhibit GABAergic neurotransmission, further promoting neuronal excitation. IL-6 can also affect synaptic plasticity, the ability of synapses to strengthen or weaken over time. This process is crucial for learning and memory, but dysregulation of synaptic plasticity can contribute to seizure activity. IL-6 can contribute to the breakdown of the blood-brain barrier, allowing immune cells and inflammatory mediators to enter the brain parenchyma. This infiltration can exacerbate neuronal dysfunction and promote seizure activity. Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) is a pleiotropic cytokine with both pro- and anti-inflammatory effects. It plays a complex role in various neurological disorders, including epilepsy and FS. In the context of FS, TNF- $\alpha$  can contribute to neuronal excitability and inflammation, but it can also exert neuroprotective effects under certain conditions. TNF- $\alpha$  can directly enhance neuronal excitability by modulating ion channels and neurotransmitter release. It can increase glutamate release and inhibit GABAergic neurotransmission, promoting neuronal excitation. TNF- $\alpha$  can induce the production of other inflammatory mediators, such as IL-1 $\beta$  and IL-6, amplifying the inflammatory response and further contributing to neuronal dysfunction. TNF- $\alpha$  can disrupt the blood-brain barrier, allowing immune cells and inflammatory mediators to enter the brain parenchyma, exacerbating neuroinflammation and promoting seizure activity. However, TNF- $\alpha$  can also exert neuroprotective effects by promoting neuronal survival and reducing excitotoxicity under certain conditions. The balance between its pro- and anti-convulsant effects likely depends on the specific context, including the timing and magnitude of its expression, the presence of other inflammatory mediators, and the specific neuronal populations involved. The elevated levels of IL-1 $\beta$ , IL-6, and TNF- $\alpha$

observed in our study suggest a complex interplay of these cytokines in the pathogenesis of FS. These cytokines do not act in isolation, they interact with each other and with other inflammatory mediators, creating a complex network of signaling pathways that influence neuronal excitability and seizure susceptibility. For instance, IL-1 $\beta$  can induce the production of IL-6 and TNF- $\alpha$ , further amplifying the inflammatory response. TNF- $\alpha$  can also stimulate the production of IL-1 $\beta$  and IL-6, creating a positive feedback loop that perpetuates inflammation. These interactions highlight the intricate nature of the inflammatory response in FS and underscore the need for a holistic approach to understanding and managing this condition. The dysregulated inflammatory response observed in children with FS may be triggered by the underlying infection or fever, or it may reflect an inherent predisposition to exaggerated inflammatory responses in certain individuals. Genetic factors, such as polymorphisms in cytokine genes or genes regulating immune responses, may contribute to this predisposition. For example, variations in the genes encoding IL-1 $\beta$ , IL-6, TNF- $\alpha$ , or their receptors could influence cytokine production or signaling, leading to an exaggerated inflammatory response. Similarly, polymorphisms in genes regulating immune responses, such as toll-like receptors (TLRs) or NOD-like receptors (NLRs), could affect the recognition of pathogens and the initiation of the inflammatory cascade. Environmental factors, such as exposure to infections, stress, or nutritional deficiencies, may also influence the inflammatory response and contribute to FS susceptibility. For instance, early-life infections have been shown to alter immune development and increase the risk of febrile seizures.<sup>11-14</sup>

Our study reveals a significant reduction in serum levels of the anti-inflammatory cytokine IL-10 in children with febrile seizures (FS). This finding adds another layer of complexity to the understanding of FS pathogenesis, highlighting the crucial role of anti-inflammatory mechanisms in maintaining immune homeostasis and neuronal balance. IL-10, a potent immunoregulatory cytokine, plays a pivotal role in dampening inflammation and resolving the immune

response. Its deficiency in children with FS suggests an impaired ability to control inflammation, potentially contributing to the unchecked pro-inflammatory state and the heightened neuronal excitability associated with this condition. To fully grasp the implications of this finding, it's essential to delve deeper into the multifaceted roles of IL-10 and explore the potential consequences of its deficiency in the context of FS. Interleukin-10 (IL-10) is a cytokine with diverse immunoregulatory functions. Produced primarily by immune cells, including macrophages, T cells, and B cells, IL-10 acts as a brake on the immune system, preventing excessive inflammation and promoting the resolution of immune responses. This regulatory function is crucial for maintaining immune homeostasis and preventing collateral damage to host tissues. IL-10 inhibits the production of pro-inflammatory cytokines, such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , by immune cells. This suppression helps to dampen the inflammatory response and prevent excessive immune activation. IL-10 can inhibit the ability of antigen-presenting cells, such as macrophages and dendritic cells, to present antigens to T cells. This inhibition reduces T cell activation and proliferation, further dampening the immune response. IL-10 promotes the differentiation of regulatory T cells (Tregs), a specialized subset of T cells that suppress immune responses and maintain self-tolerance. Tregs play a crucial role in preventing autoimmune diseases and resolving inflammation. IL-10 can directly suppress the function of effector T cells, such as Th1 and Th17 cells, which are involved in cell-mediated immunity and inflammation. The decreased levels of IL-10 observed in our study suggest an impaired anti-inflammatory response in children with FS. Without adequate IL-10, the production of pro-inflammatory cytokines, such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , may go unchecked, leading to excessive inflammation and neuronal damage. This uncontrolled inflammation can contribute to seizure activity by increasing neuronal excitability, disrupting the blood-brain barrier, and altering neurotransmitter balance. IL-10 deficiency may impair the development and function of Tregs, compromising their ability to suppress excessive immune activation and resolve inflammation. This

impairment can further contribute to the unchecked pro-inflammatory state and increase susceptibility to seizures. IL-10 has been shown to exert neuroprotective effects by reducing excitotoxicity and promoting neuronal survival. Its deficiency may exacerbate neuronal damage caused by inflammation and oxidative stress, further contributing to seizure activity. The findings of our study highlight the importance of the IL-10 axis in the pathogenesis of FS. The balance between pro-inflammatory cytokines, such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , and the anti-inflammatory cytokine IL-10 appears to be crucial for maintaining immune homeostasis and neuronal balance. A deficiency in IL-10 may disrupt this balance, tipping the scales towards excessive inflammation and increasing susceptibility to seizures. This imbalance may be further exacerbated by the interplay between inflammation and oxidative stress. Pro-inflammatory cytokines can stimulate ROS production, while oxidative stress can activate inflammatory pathways, creating a vicious cycle that amplifies both processes. IL-10, through its anti-inflammatory and antioxidant effects, can help to break this cycle and restore balance. Its deficiency may perpetuate this vicious cycle, further contributing to neuronal dysfunction and seizure activity. The impaired anti-inflammatory response observed in children with FS may be influenced by both genetic and environmental factors. Genetic variations in the IL-10 gene or in genes regulating IL-10 production may contribute to reduced IL-10 levels and impaired anti-inflammatory function. For example, polymorphisms in the promoter region of the IL-10 gene have been associated with altered IL-10 production and susceptibility to various inflammatory diseases. Similarly, variations in genes encoding transcription factors that regulate IL-10 expression, such as STAT3 and GATA3, could influence IL-10 levels and immune responses. Environmental factors, such as exposure to infections, stress, or nutritional deficiencies, may also influence IL-10 production and contribute to impaired anti-inflammatory function. For instance, early-life infections have been shown to alter immune development and affect cytokine production, potentially influencing IL-10 levels and



susceptibility to FS. The findings of our study raise the possibility of targeting the IL-10 axis for therapeutic intervention in FS. Strategies aimed at boosting IL-10 production or enhancing its anti-inflammatory effects may prove beneficial in restoring immune balance and reducing seizure susceptibility. Administering recombinant IL-10 or IL-10 analogs could potentially compensate for the deficiency and restore anti-inflammatory function. However, the safety and efficacy of this approach need to be carefully evaluated in clinical trials. Stimulating endogenous IL-10 production through pharmacological or immunological interventions could be another therapeutic strategy. For example, certain probiotics or prebiotics have been shown to enhance IL-10 production and modulate immune responses. Gene therapy approaches aimed at correcting genetic defects in the IL-10 gene or in genes regulating IL-10 production could potentially restore IL-10 levels and improve anti-inflammatory function. However, this approach is still in its early stages of development and faces significant challenges.<sup>15-17</sup>

Our study reveals a compelling link between oxidative stress and febrile seizures (FS), adding a crucial dimension to the understanding of this common childhood neurological disorder. The significantly elevated levels of malondialdehyde (MDA), a marker of lipid peroxidation, coupled with the reduced activity of superoxide dismutase (SOD), a key antioxidant enzyme, paint a clear picture of increased oxidative damage and a compromised antioxidant defense system in children with FS. This observation aligns with a growing body of research highlighting the role of oxidative stress in various neurological conditions, including epilepsy and seizures. To fully appreciate the implications of these findings, it's essential to delve deeper into the mechanisms by which oxidative stress contributes to FS pathogenesis. This exploration will encompass the sources of reactive oxygen species (ROS) in the context of FS, the detrimental effects of ROS on neuronal function, the intricate interplay between inflammation and oxidative stress, and the potential therapeutic implications of targeting oxidative stress pathways. ROS are highly reactive molecules that can damage cellular

components, including lipids, proteins, and DNA. Under normal physiological conditions, ROS are generated as byproducts of cellular metabolism, primarily within the mitochondria, the energy powerhouses of cells. However, during periods of stress, such as infection or fever, ROS production can increase dramatically, overwhelming the body's antioxidant defenses and leading to oxidative stress. Fever can disrupt mitochondrial function, leading to increased electron leakage from the electron transport chain and enhanced ROS generation. This mitochondrial dysfunction can further compromise neuronal energy production and exacerbate neuronal damage. Inflammation, a hallmark of FS, can also trigger ROS production. Activated immune cells, such as neutrophils and macrophages, release ROS as part of their antimicrobial defense mechanisms. However, excessive ROS production can damage surrounding tissues, including neurons. The metabolism of certain neurotransmitters, such as glutamate, can also generate ROS. Excessive glutamate release, which can occur during seizures, can lead to excitotoxicity and increased ROS production, further contributing to neuronal damage. Disruption of the blood-brain barrier, which can occur during inflammation, can allow the influx of ROS from the periphery, further contributing to the oxidative stress burden in the brain. ROS can inflict widespread damage on cellular components, particularly affecting vulnerable neurons. ROS can attack polyunsaturated fatty acids in cell membranes, leading to lipid peroxidation and membrane damage. This damage can disrupt neuronal signaling and impair cellular function. MDA, a byproduct of lipid peroxidation, serves as a marker of oxidative damage to lipids. ROS can oxidize amino acid residues in proteins, altering their structure and function. This oxidation can lead to protein misfolding, aggregation, and dysfunction, contributing to neuronal damage. ROS can damage DNA, leading to mutations and genomic instability. This damage can disrupt gene expression and impair cellular function, potentially contributing to neuronal death. Mitochondria, the primary source of cellular energy, are particularly susceptible to oxidative damage. ROS can damage mitochondrial DNA, proteins, and lipids,

impairing energy production and further compromising neuronal function. ROS can exacerbate excitotoxicity, a process where excessive glutamate release leads to neuronal damage and death. ROS can enhance glutamate release and impair glutamate reuptake, further contributing to excitotoxicity. The interplay between inflammation and oxidative stress is a crucial aspect of FS pathogenesis. These two processes are not independent entities, they interact and amplify each other, creating a vicious cycle that perpetuates neuronal dysfunction and seizure activity. Inflammation can trigger ROS production through several mechanisms. Activated immune cells, such as neutrophils and macrophages, release ROS as part of their antimicrobial defense mechanisms. However, excessive ROS production can damage surrounding tissues, including neurons. Pro-inflammatory cytokines, such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , can stimulate ROS production by various cells, including neurons and glial cells. Inflammation can disrupt mitochondrial function, leading to increased ROS production and further exacerbating oxidative stress. Conversely, oxidative stress can activate inflammatory pathways. ROS can activate nuclear factor- $\kappa$ B (NF- $\kappa$ B), a transcription factor that regulates the expression of various pro-inflammatory genes. NF- $\kappa$ B activation can lead to increased production of pro-inflammatory cytokines, further amplifying the inflammatory response. ROS can activate inflammasomes, multiprotein complexes that trigger the release of pro-inflammatory cytokines, such as IL-1 $\beta$  and IL-18. Inflammasome activation can further exacerbate inflammation and contribute to neuronal damage. This intricate interplay between inflammation and oxidative stress creates a self-perpetuating cycle that contributes to the pathogenesis of FS. Breaking this cycle is crucial for mitigating neuronal damage and reducing seizure susceptibility. The body has evolved a complex network of antioxidant defense mechanisms to counteract the harmful effects of ROS. Enzymatic antioxidants, such as SOD, catalase, and glutathione peroxidase, catalyze the detoxification of ROS. SOD converts superoxide radicals to hydrogen peroxide, which is then further detoxified by catalase and glutathione peroxidase. Non-enzymatic

antioxidants, such as vitamin C, vitamin E, and glutathione, scavenge ROS and prevent oxidative damage. These antioxidants act by donating electrons to ROS, neutralizing their reactivity. In the context of FS, the reduced SOD activity observed in our study suggests a compromised antioxidant defense system. This impairment may allow ROS to accumulate and inflict damage on neuronal components, contributing to seizure activity. The findings of our study highlight the importance of considering oxidative stress as a potential therapeutic target in FS. Strategies aimed at boosting antioxidant defenses or reducing ROS production may prove beneficial in mitigating neuronal damage and potentially reducing the risk of seizures. Supplementing with antioxidants, such as vitamin C, vitamin E, or coenzyme Q10, could potentially enhance antioxidant defenses and reduce oxidative damage. However, the efficacy of this approach in FS needs to be further investigated in clinical trials. A diet rich in fruits and vegetables, which are abundant in natural antioxidants, may help to boost antioxidant defenses and reduce oxidative stress. Certain drugs, such as N-acetylcysteine and melatonin, have antioxidant properties and could potentially be used to modulate oxidative stress in FS. However, further research is needed to evaluate their safety and efficacy in this context. Regular exercise and stress reduction techniques, such as meditation and yoga, have been shown to reduce oxidative stress and may prove beneficial in FS management.<sup>18-20</sup>

#### **4. Conclusion**

This study provides compelling evidence for the involvement of both inflammatory cytokines and oxidative stress in the pathogenesis of febrile seizures (FS) in children from Mumbai, India. Our findings reveal a distinct profile in children with FS, characterized by elevated pro-inflammatory cytokines (IL-1 $\beta$ , IL-6, and TNF- $\alpha$ ), decreased levels of the anti-inflammatory cytokine IL-10, and an imbalance in oxidative stress markers (increased MDA and decreased SOD). These results suggest a dysregulated inflammatory response and a compromised antioxidant defense system, potentially contributing to neuronal hyperexcitability and seizure activity. Our

study highlights the need for a holistic approach to understanding and managing FS, considering both inflammation and oxidative stress pathways.

## 5. References

1. Huang W-X, Yu F, Sanchez RM, Liu Y-Q, Min J-W, Hu J-J, et al. TRPV1 promotes repetitive febrile seizures by pro-inflammatory cytokines in immature brain. *Brain Behav Immun.* 2015; 48: 68–77.
2. Zhuang F, Zhou X, Li H, Yang X, Dong Z, Zhou W, et al. Hydrogen sulfide promotes learning and memory and suppresses proinflammatory cytokines in repetitive febrile seizures. *Neuroimmunomodulation.* 2016; 23(5–6): 271–7.
3. Kim K, Kwak BO, Kwon A, Ha J, Kim S-J, Bae SW, et al. Analysis of plasma multiplex cytokines and increased level of IL-10 and IL-1Ra cytokines in febrile seizures. *J Neuroinflammation.* 2017; 14(1): 200.
4. Kim K, Kwak BO, Kwon A, Ha J, Kim S-J, Bae SW, et al. Correction to: Analysis of plasma multiplex cytokines and increased level of IL-10 and IL-1Ra cytokines in febrile seizures. *J Neuroinflammation.* 2017; 14(1): 226.
5. Mkhize NVP, Qulu L, Mabandla MV. The effect of quercetin on pro- and anti-inflammatory cytokines in a prenatally stressed rat model of febrile seizures. *J Exp Neurosci.* 2017; 11: 1179069517704668.
6. Yamanaka G, Ishida Y, Morichi S, Morishita N, Takeshita M, Tomomi U, et al. Spinal fluid cytokine levels and single-photon emission computed tomography findings in complex febrile seizures. *J Child Neurol.* 2018; 33(6): 417–21.
7. Chen R, Li S, Wang X, Zhou J, Lu Y, Kang A. Analysis of cytokines and trace elements in children with febrile seizures. *Transl Pediatr.* 2020; 9(6): 809–17.
8. Scantlebury MH, Reid AY, Pittman QJ. Contributions of cytokines to febrile seizures. In: *Febrile Seizures.* Elsevier. 2023; 179–93.
9. Tomioka K, Nishiyama M, Tokumoto S, Yamaguchi H, Aoki K, Seino Y, et al. Time course of serum cytokine level changes within 72 h after onset in children with acute encephalopathy and febrile seizures. *BMC Neurol.* 2023; 23(1): 7.
10. Vishal V, Sachdeva A, Bhalla K, Dureja A, Tarte S. Exploring the involvement of cytokines in pediatric patients afflicted by simple febrile seizures: a case-control study. *Cureus.* 2023; 15(10): e48083.
11. Hautala MK, Helander HM, Pokka TM-L, Koskela UV, Rantala HMJ, Uhari MK, et al. Recurrent febrile seizures and serum cytokines: a controlled follow-up study. *Pediatr Res.* 2023; 93(6): 1574–81.
12. Elmas B, Erel Ö, Ersavaş D, Yürümez Y. Thiol/disulfide homeostasis as a novel indicator of oxidative stress in children with simple febrile seizures. *Neurol Sci.* 2017; 38(11): 1969–75.
13. El-Masry HMA, Sadek AA, Hassan MH, Ameen HH, Ahmed HA. Metabolic profile of oxidative stress and trace elements in febrile seizures among children. *Metab Brain Dis.* 2018; 33(5): 1509–15.
14. León Navarro DA, Crespo M, Martín M. Oxidative stress in epileptogenesis: Febrile seizures, chemoconvulsant pilocarpine, and electrical stimulation. In: *Oxidative Stress and Dietary Antioxidants in Neurological Diseases.* Elsevier. 2020; 81–94.
15. Polat I, Karaoglu P, Ayanoglu M, Cirali C, Bayram E, Yis U, et al. Inflammation and anemia in simple febrile seizures and complex febrile seizures. *Ann Med Res.* 2021; 28(10): 1835.
16. Tang L, Chen J-R. The predictive value of hemocytometry based on peripheral platelet-related parameters in identifying the causes of febrile seizures. *J Inflamm Res.* 2021; 14: 5381–92.
17. Ikeda-Matsuo Y, Tomori N, Takemoto Y, Uematsu S, Akira S, Takahashi T. Role of microsomal prostaglandin E synthase-1 in

hippocampal inflammation after repetitive febrile seizures in mouse pups. Proceedings for Annual Meeting of The Japanese Pharmacological Society. 2022; 96(0): 3-B-007-2.

18. Çetin İD, Çetin O. A preliminary study on the association between prognostic nutritional index and neutrophil-to-lymphocyte ratio with nutritional status and inflammation in febrile children's susceptibility to seizures. Rev Assoc Med Bras. 2022; 70(7): e20240166.
19. Barrett KT, Roy A, Ebdalla A, Pittman QJ, Wilson RJA, Scantlebury MH. The impact of inflammation on thermal hyperpnea: Relevance for heat stress and febrile seizures. Am J Respir Cell Mol Biol. 2021; 71(2): 195–206.
20. Söğütlü Y, Altaş U. Predictive value of neutrophil-lymphocyte ratio and other inflammation indices in febrile seizures in children. J Clin Med. 2022; 13(17): 5330.